

Article

Planning Landscape Corridors in Ecological Infrastructure Using Least-Cost Path Methods Based on the Value of Ecosystem Services

Jung A. Lee ^{1,2}, Jinhyung Chon ^{3,*} and Changwoo Ahn ⁴

¹ Department of Environmental Science and Policy, George Mason University, 4400 University Drive, MS 5F2, Fairfax, VA 22030, USA; E-Mail: arch-jung@korea.ac.kr

² Institute of Life Science and Natural Resources, College of Life Sciences and Biotechnology, Korea University, 145 Anamro, Seongbuk-gu, Seoul 136-713, Korea

³ Division of Environmental Science and Ecological Engineering, College of Life Sciences and Biotechnology, Korea University, 145 Anamro, Seongbuk-gu, Seoul 136-713, Korea

⁴ Department of Environmental Science and Policy, George Mason University, 4400 University Drive, MS 5F2, Fairfax, VA 22030, USA; E-Mail: cahn@gmu.edu

* Author to whom correspondence should be addressed; E-Mail: jchon@korea.ac.kr; Tel.: +82-2-3290-3048; Fax: +82-2-3290-3624.

External Editor: Marc A. Rosen

Received: 31 July 2014; in revised form: 17 October 2014 / Accepted: 17 October 2014 /

Published: 28 October 2014

Abstract: Ecosystem service values have rarely been incorporated in the process of planning ecological infrastructure for urban areas. Urban ecological infrastructure is a network system of natural lands and waters that provides ecosystem services. The purpose of this study was to design landscape corridors that maximize the value of ecosystem services in ecological infrastructure planning. We explored the optimal corridors to enhance the connectivity among landscape elements to design an ecological infrastructure for the city of Gwacheon, South Korea, as an example of a small urban area. We calculated the value of ecosystem services using standardized estimation indices based on an intensive review of the relevant literature and employed the least-cost path method to optimize the connectivity of landscape structural elements. The land use type in the city with the highest estimated value of ecosystem services was the riparian zone (*i.e.*, 2011 US\$7,312.16/ha). Given areal coverage of all land use types, the estimated value of developed area open spaces was 2011 US\$899,803.25, corresponding to the highest contribution to the total value of ecosystem

services. Therefore, the optimal configured dispersal corridors for wildlife were found from the riparian zones (source area) to the developed area open spaces (destination area) in the city. Several challenges remain for improving the estimation of the value of ecosystem services and incorporating these ecosystems in ecological infrastructure planning. Nonetheless, the approaches taken to estimate the value of ecosystem services and design landscape corridors in this study may be of value to future efforts in urban ecological infrastructure planning.

Keywords: ecosystem service; value of ecosystem services; least-cost path method; ecological infrastructure; landscape connectivity; landscape structure elements

1. Introduction

The term “ecosystem services” refers to the benefits gained from the complex interactions between the human environment and the functions of an ecosystem, such as carbon sequestration, improvements in air and water quality, microclimate regulation, biodiversity and recreational and cultural and social value [1–6]. Ecosystem services are considered to be a key factor for leading a healthy and prosperous life, and they are significant for the sustainability of human society [7,8]. Balmfor *et al.* [4] suggest that ecosystem services are based on basic ecosystem functions that are crucial for supporting services. Ecosystem functions substantiate ecosystem services, such as the provision of harvested crops, landscape aesthetics and habitats for biodiversity and the regulation of environmental quality [9]. In other words, ecosystem functions represent the capacity of an ecosystem to provide goods and services to indirectly satisfy human needs [10,11]. These ecosystem functions act at the intersection between ecosystem features (e.g., built-up areas and residential functions or forest and timber production) and structures (e.g., landscape diversity, complexity and fragmentation). Ecosystem structures or patterns can affect landscape functions through the composition and configuration of landscape elements [9,12,13], and they can affect ecosystem services indirectly. This statement implies that ecosystem services are expected to fulfill many landscape functions according to the landscape structure. This concept of ecosystem services has become significant among researchers and practitioners in landscape planning and management [14].

The importance of considering ecosystem services when studying landscape planning has been emphasized [14–20]. Previous studies have focused on successful ecological corridor planning in terms of conduits for wildlife [21]. However, simple ecological corridors still lack alternative strategies for addressing the ecological impact of fragmentation and the concept of ecological networks [22]. This concept has recently developed as a popular concept in urban planning, which can be seen as preserving wildlife habitats, as well as contributing to the urban ecosystem’s health and resilience. According to Rapport *et al.* [23], the provision of ecosystem services is essential for a healthy urban ecosystem. The ecological networks are necessary to develop approaches that integrate biophysical, economic and sociocultural effects from ecosystem services, in order to propose a better way of landscape planning and management in urban areas.

As rapid urbanization continues, it is assumed that more than 60% of the world’s population will live in urban areas by 2030 [24]. With this increased concentration, urban infrastructure has been intensively

developed, with little linkage to natural ecosystems [20]. This trend has led to the fragmentation and isolation of urban green spaces, negatively affecting potential ecosystem services [25,26]. There are several ecosystem services available in urban areas [20,24–27] that produce positive biophysical, economic and sociocultural effects. The demand on ecosystem services has continued to increase with population growth [28,29]. Many efforts to factor ecosystem services into landscape planning have been made in the past several decades [14–18]. Recent research shows that an integrated framework, such as ecological infrastructure, is necessary to achieve a systematic and comprehensive assessment of the value of ecosystem services for setting policies and decision making in landscape planning [14–17]. Ecological infrastructure is defined as a network system of natural lands that provides ecosystem services [19]. The development of ecological infrastructure has recently become a practical strategy for providing or supporting ecosystem services to maximize the benefits of the ecosystem. Wetlands are particular practical ecological infrastructures of urban development in the ecological engineering field. Many studies have comprehensively and intensively dealt with constructed wetlands through ecological valuation schemes and a set of indices to evaluate ecosystem services, including the environmental impact and emissions [30–36]. In the field of landscape planning and management, the main challenge has become determining the optimal allocation and management of several different land use types through ecological infrastructure planning [14]. In other words, the large-scale ecological network, which is created by the connection between small or large ecological infrastructure, such as wetlands, has become a major issue of ecological infrastructure in urban areas. This study is also a part of this paradigm.

The design of ecological infrastructure is typically based on connections between “patches” and “matrices” via “corridors”, representing the main spatial landscape structure elements of the landscape network [37]. The least-cost path method has been shown to be useful in several studies [38–42] for determining paths to connect landscape structure elements, which are commonly integrated, for example, in geographic information system (GIS) technology. Least-cost path analysis is one of the best methods for achieving the optimal establishment of paths between landscape elements (e.g., large hubs (matrices), smaller sites (patches) and links (corridors)) [43]. This useful method is based on the fact that paths for wildlife movement are affected by the characteristics of landscape, including the land cover, roads and slope [41,44]. Each cell of a raster dataset is allocated a value according to the cost of movement. A value that would incorporate the estimated value of ecosystem services can be assigned based on land cover characteristics, which is easily realized within the raster data cells used for the study. The value of ecosystem services often depends on the maintenance of biodiversity [45,46]. As wildlife species are affected differently by land cover characteristics, including impervious surface and natural green surfaces, the weights of land cover for wildlife habitats contribute to biodiversity [47–49]. The weights of land cover are based on the value of ecosystem services using the least-cost path method. The least-cost path method creates the best travel path according to the composition of the cells that are assigned the shortest distance with the least resistance between two patches of wildlife habitats [38]. Thus, this method can help generate the best theoretical path to connect suitable habitats for wildlife dispersal. In several studies, the least-cost path method has been employed for the analysis of landscape connectivity based on considering wildlife dispersal, which is faster and more convenient for the visualization of landscape connectivity than any other method [50–52], such as random walk modeling [53–55], network analysis [56,57] or gravity models [58]. This approach can also be used to maximize ecosystem services

by connecting landscape structure elements in designing corridors for ecological infrastructure. The landscape structure elements may be influenced by ecological processes related to ecosystem services [59].

The purpose of this study is to design landscape corridors that maximize the value of ecosystem services in ecological infrastructure planning. These corridors are meant to serve as connection paths between the main wildlife habitats in the city of Gwacheon. The city of Gwacheon, located in the vicinity of Seoul, South Korea, was chosen for the study. The specific objectives were (1) to calculate the least cost based specifically on ecosystem service values using a least-cost path model and (2) to design a framework demonstrating corridors that serve as the best connection paths between suitable habitats in the city.

2. Methods


2.1. Study Area

The city of Gwacheon, located in the mid-western region of Gyeonggi-do near Seoul, South Korea, is a medium-scale urban area with considerable green space. The study site is geographically situated at 37°23'53"N to 37°27'52" and 127°02'52"E to 126°57'52". The site encompasses an area of approximately 35.86 km² and includes 32.45 km² of natural areas, such as forest. The climate of Gwacheon is characterized by high temperatures and high humidity in summer and cold, dry conditions in the winter.

The landscape of this area has various characteristics in terms of natural resources and cultural attractions. The city is located in a basin surrounded by mountains (Mt. Gwanak, Mt. Cheonggyes and Mt. Umyeon) and includes several streams (Yangjaecheon, Makgyecheon and Galhyeoncheon). The population is distributed along the rivers. Similar to most urban areas, however, it has experienced a rise in population, technology and infrastructure, such as the development of the Government Building and highways connecting the region with Seoul and other surrounding cities. Figure 1 shows the land use pattern in Gwacheon in a land cover map. The urban ecosystems of the city have become highly modified and fragmented.

2.2. Geospatial Data

We collected a geospatial dataset to represent the landscape of Gwacheon by downloading land cover data from the Ministry of Environment and digital topographic maps from the Ministry of Land, Infrastructure and Transport. The current land cover map (2000, with a scale of 1/25,000) and a map of major rivers and traffic districts (2008, with a scale of 1/25,000) were interpreted and modified from aerial photographs (2009). To effectively assign a generalized value of ecosystem services for the study site, the land use types within the study site were reclassified into 10 primary land use types: developed areas of low and medium intensity, developed areas of high intensity, developed area open spaces, rice paddies, croplands, orchards, forests, grasslands, riparian zones and bare soils. Highway and road information was obtained from line data from a topographic map (2007, with a scale of 1/25,000) and aerial photographs. Using a digital topographic map, 30-m digital elevation model (DEM) data were generated from contour lines. The slope was calculated from the 30-m DEM data. The Arcmap 10.1 GIS platform [60] was used for the digitization and analysis of the dataset, then converted to rasters as a 30 × 30-m² grid.

Figure 1. Land cover map of Gwacheon.

2.3. Planning Landscape Corridors in the Ecological Infrastructure

This study, aimed at planning landscape corridors, was based on least-cost path methods, where the value of ecosystem services was assessed to determine the best connection among wildlife habitats and to maximize the ecosystem services in Gwacheon. The value of the ecosystem services for each land cover type present in Gwacheon was estimated using synthesis and standardization coefficients for the value of ecosystem services from previous research. We developed least-cost paths for ecological infrastructure planning according to the estimated value of ecosystem services, which were created based on the shortest distance with the least resistance to the movement of the main species in Gwacheon. The resistance encompassed the value of ecosystem services for each land cover type, the density of roads/highways and slopes.

2.3.1. Estimated Value of Ecosystem Services

The value of ecosystem services generally represents the creditable nonmarket value based on efficiency and cost-effectiveness values, including the degree of sustainable use within the complex conditions of the ecosystem and equity for the enhancement of human wellbeing [61]. Identifying and quantifying the value of ecosystem services helps us to better understand the benefits of these services and the provision of ecosystems for human society [62]. In addition, a method for identifying and quantifying the value of ecosystem services could be a valuable tool for the efficient allocation of ecosystem benefits [11,62–64]. Quantifying the value of ecosystems could provide clues about how to achieve the social recognition and acceptance of ecosystem management across multiple geographic scales [65]. The most widely used and best known estimation values for ecosystem services are a set of “generalized coefficients” proposed by Costanza *et al.* [6,66–69]. However, these coefficients are not without limits or constraints on their use [70]. Thus, many researchers made subsequent efforts, which resulted in a range of generalized coefficients that are applicable for estimating the value of ecosystem services for each land use type from varying regions [71–76]. We adopted all of the available values from the existing literature and created our own set or range of coefficients relevant to the land use types included in this study (Table 1).

Table 1. Value estimates for each land use type based on a number of studies.

Land Use Typology ^a	Definition ^b	Total of Service Means Values (2011 US\$/ha/Year)	Total SD of Means (2011 US\$/ha/Year)	Source
Developed areas of low and medium intensity	Includes areas with a mixture of constructed materials and vegetation. Impervious surfaces account for 20%–79% of the total cover.	0	0	[6,67,72]
Developed areas of high intensity	Includes highly developed areas where people reside or work in high numbers. Impervious surfaces account for 80%–100% of the total cover.	0	0	[6,67,72]

Table 1. Cont.

Land Use Typology ^a	Definition ^b	Total of Service Means Values (2011 US\$/ha/Year)	Total SD of Means (2011 US\$/ha/Year)	Source
Developed area open spaces	Includes areas with a mixture of constructed materials, but mainly contains vegetation in the form of lawn grasses. Impervious surfaces account for less than 20% of the total cover.	2130.67	1189.36	[6,77]
Rice paddies	Areas where perennial herbaceous vegetation accounts for more than 80% of the vegetative cover and the soil or substrate is periodically saturated with or covered with water, such as rice paddies.	5131.91	1540.98	[6,68,78–82]
Croplands	Areas being used for the production of crops other than rice. Plantations with cash crops, such as herbal teas or horticultural products.	413.07	162.33	[6,68,83–85]
Orchards	Areas being used for food production by planting trees and shrubs, such as <i>Malus pumila</i> Mille, <i>Pyrus serotina</i> Rehder and <i>Diospyros kaki</i> .	594.80	16.48	[6,67]
Forests	Includes natural forest plantations. Lands with tree-canopy cover account for more than 20%. The trees should be able to reach a minimum height of 5 m.	1937.03	1719.01	[6,67,78,85–103]
Grasslands	Includes infertile or degraded land where no trees or shrubs grow.	315.19	65.96	[6,67,68,83]
Riparian zones	All areas of open water, including riparian buffer zones, generally with less than 25% vegetation or soil cover	7312.16	5836.06	[6,68,91,104–110]
Bare soils	Areas of bedrock, desert pavement, scarps, talus, slides, volcanic material, glacial debris, sand dunes, strip mines, gravel pits and other accumulations of earthen material. Vegetation generally accounts for less than 15% of the total cover.	25.98	0.00	[6,68]

^a The land unit typologies consisted of the land cover types present in Gwacheon; ^b source: [111].

Peer-reviewed articles from 15 journals were collected, synthesized and combined to develop a well-defined estimated value of ecosystem services. All of the potential data on such values were screened using the keywords “ecosystem service”, “value of ecosystem service” and “ecosystem services value”. Data were extracted from studies that provided a monetary value for a given ecosystem service attached to a specific land use type and a specific time period. The values of ecosystem services were applied to enable the conversion of monetary values to per-hectare values. Because the values of

ecosystem services have been reported in many different metrics and currencies for different time periods, locations and price levels, it was necessary to organize these values in a standardized form to reduce generation errors. We developed estimation indices for the existing values of ecosystem services based on this literature review to find well-defined coefficients. The estimation indices entailed obtaining economic estimates for the value of nonmarket services that are not directly traded in markets for environmental resources [75]. Several of the coefficients used to estimate the value of ecosystem services exhibited a central tendency [71,112,113].

In the review of 37 studies, the estimated values of ecosystem services were taken as averages of the existing estimated values (Table 1). The average of all collected values was estimated using a range of approaches [78]. To aid in the direct comparison and aggregation, the collected ecosystem service values were standardized to common spatial, temporal and currency units, namely, international dollars per hectare per year (\$/ha/year) for 2011 (Table 1). When necessary, the estimates were converted into the official local currency. The values were then adjusted to 2011 values using the nominal GDP per capita of each country. The official exchange rates and nominal GDP per capita were determined according to MeasuringWorth.com [114]. The sum of all ecosystem service values constituted the total service mean values (Table 1). The total of service mean values included three types of ecosystem services (regulating, provisioning and cultural services), and the three types of ecosystem services were composed of 10 ecosystem functions (gas and climate regulation, disturbance regulation, water regulation, waste treatment, water supply, food production, pollination, refugia, recreation and cultural) [6,20] (Table 1).

This study used the assessment method developed by [66] to calculate the value of ecosystem services for each land cover type. The formula for calculating the values is as follows:

$$[\text{Vaue of ecosystem services} = \sum(A_k \times V_k)] \quad (1)$$

where A_k is the area (ha) of the corresponding cover type, V is the annual value of the land cover per hectare and k is the land cover type. Each value for each ecosystem service category was estimated by calculating the estimated value for each land use type.

2.3.2. Least-Cost Path Method

The least-cost path method using GIS-based analysis supported the best corridors for connecting landscape patches to be adopted in ecological infrastructure planning for Gwacheon. The least-cost path method is based on cost-distance analysis. Two GIS layers (the source layer and the resistance layer) are used as inputs in the least-cost path method. The source layer represents the landscape patches from which connectivity is calculated according to the cost benefits of ecosystem services related to changing ecological infrastructure into urban infrastructure. The resistance layer represents resistance values assigned by summarizing the weight values of the relative costs of traveling through each land use type and geographical position. The weight values are formed by each cell (30 m × 30 m) in the grid based on the value of ecosystem services according to the land use-type attributes. In addition, the relative cost of traveling is based on the consideration of the main wildlife in Gwacheon, which includes medium-large-sized mammals, such as *Hydropotes inermis* and *Prionailurus bengalensis*, and their movement according to the land use type and geographical position, including the density of roads/highways and slopes. To increase connectivity between landscape patches based on the

composition and configuration of the landscape structure, a weight value for each land use type was inversely calculated in proportion to the lowest ecosystem service value for a land use type according to the value of ecosystem services for each land use type. The weights were calculated as follows, where w is the computed weight of ecosystem services for a land use type:

$$[w_i = \frac{\sum_{i=1}^n a_i - a_i}{\sum_{i=1}^n a_i} \times 100, \text{ for all values of ecosystem services } i = 1, 2, \dots, n] \quad (2)$$

The weight value for the geographical position was set to one within a cell, which reflects a minimum cost movement [115,116]. The resistance value was ultimately assigned to the cost-surface raster by summarizing the relative weight cost of each land cover using the Calculation Cost Surface Tool in ArcToolbox. The resistance value for each cell indicates the path to the source, measured as the least cost involved in moving over the resistance layer.

There are four steps in the least-cost path method to find the best paths for ecological infrastructure planning: assignment of the source area; creation of a cost-surface raster; assignment of the destination area; and creation of potential least-cost paths between each source/destination pair. The source area and destination were assigned to calculate and generate a cost-weighted surface. The cost-weighted surface is the cost value related to movement in each grid, which can be generated through superposition of the cost raster, source area and destination. The geospatial data included the land cover map, comprising the land use type and geographical position in Gwacheon. In addition, all of the values of ecosystem services for different land use types were estimated to create a cost-surface raster. All of the geospatial data were analyzed using Arcmap 10.1 [61] for each 90 m² grid.

3. Results and Discussion

3.1. Ecosystem Service Estimates for the City of Gwacheon

Table 1 shows the estimated ecosystem service values calculated per land use unit, garnered from a number of studies. The total mean service values were calculated from each value of ecosystem services within each land use unit typology (Table 1). The statistical means were calculated with medians from the minimum, maximum or single values available in the literature, summed as the total mean value for all studied ecosystem services. The total standard deviation shows the range of the total mean service values.

The developed areas of low and medium intensity and developed areas of high intensity did not present any anticipated ecosystem service value. These land use types include residential (mainly suburban) and commercial (mainly urban) areas where paving materials are present, without biomes. Most land use types based on natural resources, such as forests, water and vegetation, show a capacity to provide ecosystem services. For the developed area open spaces, recreation services appear to be the main ecosystem service provided; thus, there is a need to assess more detailed values of recreation services, because this land use type is similar to natural resources as a form of ecological infrastructure at the center of the urban ecosystem.

The total value of riparian zones ranked first at \$7312.16 (Table 1). This result implies that riparian zones must be a main consideration in the ecological infrastructure of urban areas. Riparian zones are highly valuable in urban ecosystems for a range of environmental, social and economic reasons [117]. Most major cities have been built along rivers, which play roles in transportation, are related to industrial

functions and provide a natural environment for people, as well as a stable drinking water supply [118]. To protect urban riparian areas, many river commissions have been established for planning and management purposes [119]. In the same vein, riparian zones are an important landscape patch in the ecological infrastructure, because they provide many valuable ecosystem services to society.

The total value of rice paddies ranked second at \$5131.91, and developed area open spaces ranked third at \$2130.67 (Table 1). Rice paddies not only produce rice for use as food, but also provide multiple ecosystem services as artificial freshwater wetlands. According to the OECD [120], rice paddies exhibit a vast potential to supply multiple ecosystem services, including food security, maintenance of the viability of rural communities, environmental protection, sustainable management of renewable natural resources, preservation of biodiversity and aesthetic landscapes, despite also having features with negative services, such as high methane emissions. Developed area open spaces received the top ranking, because they include parks, recreation areas and small urban green areas, which could be part of the ecological infrastructure for enhancing societal wellbeing.

The total value of ecosystem services for each land use unit typology in Gwacheon is presented in Table 2. Synthesized and standardized ecosystem service values were used to calculate and estimate the total value of ecosystem services. The total value of ecosystem services was estimated to be \$5,673,026.83. The value of forests was highest (\$4,132,227.36), followed by the value of developed area open spaces (\$899,803.25). Forests are located on the edge of town in Gwacheon, occupying over 60% of the area (2133.28 ha). Because urban ecological infrastructure planning focuses on urban areas, the main patch is riparian zones, not forests, due to their geographical location. The value of not only developed area open spaces (\$899,803.25), but also riparian areas (\$297,678.04) are located in one of the most densely populated regions of Gwacheon, which has around 42% of the total population (11,069 persons) according to the 2009 population census. As a high economic value of ecosystem services depends on a high residential population [121], the value of riparian zones seems to be influenced by residential population in the city. Those values indicate that such highly populated areas show great potential to improve human wellbeing and provide benefits from ecosystem services in urban areas.

Table 2. Land use types and annual value of ecosystem services per land use type in Gwacheon.

Land Use Type	Component	Estimated Value (2011 US\$/ha)	Area (ha)	Value of Ecosystem Services (2011 US\$)
Developed areas of low and medium intensity	Residential		235.36	0.00
Developed areas of high intensity	Commercial/traffic district/parks and recreation		198.36	0.00
Developed area open spaces	Public facilities	2130.67	422.31	899,803.25
Rice paddies	Rice paddy	5131.91	34.03	174,638.90
Croplands	Farmland/greenhouse	413.27	342.06	141,363.14
Orchards	Orchard	594.80	22.39	13,317.57
Forests	Evergreen/deciduous/mixed	1937.03	2133.28	4,132,227.36
Grasslands	Natural/artificial	315.19	42.2	13,301.02
Riparian zones	Inland water	7312.16	40.71	297,678.03
Bare soils	Bare soil	25.98	26.85	697.56
Total		17861.01	3497.55	5,673,026.83

3.2. Planning Landscape Corridors within the Ecological Infrastructure


The planning of landscape corridors in the ecological infrastructure was conducted by finding the best paths for improving connectivity between landscape patches constituting suitable habitats for wildlife based on the value of ecosystem services. We created a potential dispersal corridor for continuous wildlife habitats in the urban area of Gwacheon. This corridor linked the highest value of ecosystem services per hectare, which occurred in riparian zones. The least-cost path method contributed to the search for an appropriate linkage corridor in ecological infrastructure planning. Based on the value of ecosystem services per land use type, connecting riparian zones and developed area open spaces could be a suitable strategy for maximizing ecosystem services in Gwacheon. Under these circumstances, the least-cost path method was applied in four steps. Spatial data were collected from a land cover map, and the value of ecosystem services was assessed.

First, source areas were assigned to calculate and generate a cost surface. A source area represents a large hub linking the ecological infrastructure, which in this case relates to the riparian zones, *i.e.*, Yangjaecheon and Makgyecheon (Figure 1). These riparian zones are representative of high-value ecosystem service areas among the landscape elements present in Gwacheon. Riparian zones have also been assigned as nature conservation and environmental protection areas by the Ministry of the Environment in Korea. This land use type, which includes the interface between terrestrial and aquatic components of the landscape, provides useful food resources with large amounts of nutrients to wildlife habitats.

Second, to generate a cost surface, which is the map-represented cost value associated with the movement in each grid (90 m²), this study considered three components related to the cost: land use types, roads/highways and slopes. Land use types were categorized based on the value of ecosystem services. Table 3 shows the resistance values obtained from the weight value for land use types based on the value of ecosystem services as a function of Equation (2) and the weight of the other elements (roads/highways and slopes) to create a cost-surface raster. The cost-surface raster was created according to the integration of four components (land use type, density of road, highway, slope) by summarizing the relative weight cost of each land cover. Passing the values of ecosystem service through a land use type is a significant factor in linking the ecological infrastructure to increase the total value of an ecosystem in urban planning (Table 3). Roads and highways were represented as density values, and the slope represented the resistance value related to wildlife movement. These factors are associated with the concept of landscape permeability, which evaluates landscape connectivity and describes the ability of wildlife to pass through certain environments. The cost-surface raster was generated using weights for these components referring to the literature [115,116]. To understand the concept of landscape permeability, the weight values for different land use types were calculated adversely using the value of ecosystem services in the land use types present in Gwacheon. Furthermore, the cost-surface raster can aid in the configuration of land use types according to the value of ecosystem services. A cost-surface raster map was created based on the shortest distance with the least resistance, as shown in Figure 2. This map shows the results for the distance cost of the resistance values for the source area in gray scale. The increasing cost values (darker colors) toward the riparian area, in the center of Gwacheon, demonstrate the effect of the distance from the source area (Figure 2).

Table 3. Weight per land unit, density of roads/highways and slopes for resistance values.


Component		Weight Value	
Land use type	Developed areas of low and medium intensity	100.00	0.65
	Developed areas of high intensity	100.00	
	Developed area open spaces	84.14	
	Rice paddies	96.92	
	Croplands	97.51	
	Orchards	99.77	
	Forests	27.16	
	Grasslands	99.77	
	Riparian zones	94.75	
	Bare soils	99.99	
Density of road	0–1 km/km ²	1	0.15
	1–2 km/km ²	2	
	2–4 km/km ²	5	
	4–6 km/km ²	7	
	More than 6 km/km ²	10	
Highway	Less than eight lanes	100	
	More than eight lanes	200	
Slope	$10(1.0 - (1.0/1.0 + e^{-(\text{slope}-30)/7}))$		0.2

Figure 2. Cost-surface raster map for determining the connectivity of landscape elements.

Third, destination areas were selected where there were developed area open spaces. These areas are fragmented natural habitats in an urban area (Figure 1). To increase the connectivity of each habitat, a developed area open space was selected as the destination area.

Finally, least-cost paths were created from the cost-surface raster map (Figure 3). The cost-surface raster map was obtained using the cost distance, and the direction raster began at the source area and ended at the destination. Figure 3 shows the results of all calculations connecting the source area and destination area to assess the connectivity of landscape patches in Gwacheon. A darker path color indicates a stronger landscape patch connection (Figure 3). As the least-cost path is based on the cost surface calculated in the preceding step, the least-cost path is the best potential linear dispersal corridor for improving ecosystem services. This result could be useful in the planning of ecological infrastructure to increase the value of ecosystem services in Gwacheon.

Figure 3. Best potential dispersal corridor from the source areas (riparian areas) to destination areas (developed area open spaces) in ecological infrastructure planning to enhance ecosystem services in Gwacheon. Ecological infrastructure is defined as a network system of natural lands that provides ecosystem services.



The lengths and widths of the corridors were not considered when dispersal corridors from the main habitats were occupied by species, as this planning approach is sufficiently capable of producing a conceptual plan for ecological infrastructure planning by allocating the arrangement and location of corridors to achieve landscape connectivity. Landscape connectivity arises from not only the movement of organisms, but also their spatial distribution [122,123]. The extension of the area of a land use type affects the value of the ecosystem services of that land use type, which was reflected in the least-cost path analysis using the resistance value for the shortest distance of corridors. Concerning the aspects of the spatial distribution in landscape connectivity, the value of the ecosystem services of a land use type

focused on determining the characteristics of each land use type rather than the pattern of landscape patches. The results are integral to understanding the processes involved in the interface between ecosystem functions and structures from the viewpoint of the value of ecosystem services.

The appropriate types of ecological infrastructure could be determined through assessing the value of ecosystem services in the riparian zones. Because the planning of ecological infrastructure targets the riparian zones in urban areas, the appropriate ecological infrastructure along the best corridor must be considered based on the results of the assessment of ecosystem service values. Water regulation, waste treatment and recreation values are ranked highly in riparian zones. Therefore, the selected ecological infrastructure must maximize these ecosystem service values. For example, a wetland shows excellent water regulation and waste treatment capabilities [71,88]. Greenways could also be suitable for enhancing recreation services. A greenway is a representative ecological infrastructure that provides opportunities for recreation for urban dwellers [124]. However, because ecosystem services rely on different ecological processes and spatial patterns, ecological infrastructure planning should be considered a specific ecosystem service and ecological target. In the case of riparian habitats (e.g., floodplain wetlands), the connectivity might need to be limited for the ecosystem service of biodiversity [125].

4. Conclusions

In this study, we investigated the best corridors for enhancing connectivity among landscape elements to design an ecological infrastructure for the city of Gwacheon, South Korea. The generated least-cost path provides dispersal corridors from riparian zones to developed area open spaces, which present a high ecosystem service value via providing suitable wildlife habitats. This set of corridors passes through residential areas containing more green cover, as well as through agricultural areas, such as rice paddies, croplands and orchards. Implementing corridors scattered between patches of urban areas in Gwacheon would clearly be favorable for landscape connectivity. The aim of ecological infrastructure planning is to promote ecosystem and human health in urban areas. Our planning of landscape corridors using the least-cost path method demonstrated the possibility of applying the value of ecosystem services in the ecological infrastructure, which means that the outcomes of planning contribute to human health through providing healthy wildlife habitats in urban areas.

The challenge of synthesizing the estimated values of ecosystem services was at least partially surmounted through using values from databases generated in previous studies, translated for application to the present work. However, there are a number of limitations to the applicability of these data. There is a possibility for the overestimation or underestimation of the results. In addition, the literature values did not correspond to most of the literature addressing assessments of the value of ecosystem services per land use type, focusing on ecosystem services in Europe and South America, whereas our study site was in South Korea. In addition, the quantity of available data was insufficient. Thus, estimation indices for the value of ecosystem services must be developed in consideration of the environment of South Korea. Future studies must focus on creating a database for estimation indices based on regional environmental characteristics. Nevertheless, the planning landscape corridors within ecological infrastructure planning were well supported by identifying the estimated value of ecosystem services. Our study was intended to review all possible ecosystem services to explore the opportunistic benefits of ecosystem services that may be supported by creating landscape connectivity in urban planning.

The outcome of the study may seem preliminary in that it did not relate the specific type of ecosystem service regarding the ecological corridor or the connectivity being designed in the landscape.

Our findings have the following implications for planning. First, the approach taken to estimate the value of ecosystem services in this study provided a useful example of assembling these values from a variety of spatial contexts. Second, the study represents not only an exploratory step toward a better understanding of the ecological infrastructure planning process, but also a novel landscape corridor design that supports landscape connectivity. Finally, the small urban planning case explored in this study could help create and improve policies for a sustainable urban ecosystem. In most small urban areas, local governments can promote land use policies that would provide sustainable living environments for humans.

Acknowledgments

This research was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF-2013R1A1A2058315) funded by the Ministry of Education. This research was also supported by a Korea University Grant. Additionally, thanks go to the postdoctoral scholar and affiliated faculty position provided by George Mason University while the first author was working on the manuscript. We also thank the anonymous reviewers who provided useful comments and suggestions to improve this manuscript.

Author Contributions

Jung A Lee contributed to the development of the idea and participated in all phases. Jinhyung Chon also contributed to development of the idea and helped perform the analysis with constructive discussions. Jinhyung Chon and Changwoo Ahn provided guidance for writing this paper. Changwoo Ahn helped improve the figures and manuscript. All of the authors made contributions to the work in this paper.

Conflicts of Interest

The authors declare no conflict of interest.

References

1. Schäfer, R.B. Biodiversity, ecosystem functions and services in environmental risk assessment: Introduction to the special issue. *Sci. Total Environ.* **2012**, *415*, 1–2.
2. Costanza, R.; Daly, H.E. Natural capital and sustainable development. *Conserv. Biol.* **1992**, *6*, 37–46.
3. Daly, H.E.; Farley, J. *Ecological Economics: Principles and Applications*, 2nd ed.; Island Press: Washington, DC, USA, 2010; pp. 74–75.
4. Balmfor, A.; Rodrigues, A.S.L.; Walpole, M.; ten Brink, P.; Kettunen, M.; Braat, L.; de Groot, R.S. *The Economics of Ecosystems and Biodiversity: Scoping the Science*; European Commission: Cambridge, UK, 2008; pp. 29–34.
5. Fisher, B.; Turner, R.K.; Morling, P. Defining and classifying ecosystem services for decision making. *Ecol. Econ.* **2009**, *68*, 643–653.

6. Costanza, R.; d'Arge, R.; de Groot, R.; Farber, S.; Grasso, M.; Hannon, B.; Limburg, K.; Naeem, S.; O'Neill, R.V.; Paruelo, J. The value of the world's ecosystem services and natural capital. *Nature* **1997**, *387*, 253–260.
7. Van Wijnen, H.J.; Rutgers, M.; Schouten, A.J.; Mulder, C.; de Zwart, D.; Breure, A.M. How to calculate the spatial distribution of ecosystem services: Natural attenuation as example from The Netherlands. *Sci. Total Environ.* **2012**, *415*, 49–55.
8. Daily, G.C. In *Nature's Services: Societal Dependence on Natural Ecosystems*; Daily, G.C., Ed.; Island Press: Washington, DC, USA, 1997; pp. 1–10.
9. Willemsen, L.; Verburg, P.H.; Hein, L.; van Mensvoort, M.E.F. Spatial characterization of landscape functions. *Landsc. Urban Plan.* **2008**, *88*, 34–43.
10. De Groot, R.S. *Functions of Nature, Evaluation of Nature in Environmental Planning, Management and Decision Making*; Wolters-Noordhoff: Groningen, The Netherlands, 1992; pp. 340–345.
11. Millennium Ecosystem Assessment. *Ecosystems and Human Well-Being: A Framework for Assessment*; Island Press: Washington, DC, USA, 2003; pp. 26–84.
12. Antrop, M. Background concepts for integrated landscape analysis. *Agric. Ecosyst. Environ.* **2000**, *77*, 17–28.
13. MacGarigal, K.; Marks, B. *FRAGSTATS: Spatial Pattern Analysis Program for Quantifying Landscape Structure*; U.S. Department of Agriculture: Portland, OR, USA, 1995; pp. 12–20.
14. De Groot, R.S.; Alkemade, R.; Braat, L.; Hein, L.; Willemsen, L. Challenges in integrating the concept of ecosystem services and values in landscape planning, management and decision making. *Ecol. Complex.* **2010**, *7*, 260–272.
15. Chan, K.M.A.; Shaw, M.R.; Cameron, D.R.; Underwood, E.C.; Daily, G.C. Conservation planning for ecosystem services. *PLoS Biol.* **2006**, *4*, 2138–2152.
16. Gret-Regamey, A.; Walz, A.; Bebi, P. Valuing ecosystem services for sustainable landscape planning in Alpine regions. *Mt. Res. Dev.* **2008**, *28*, 156–165.
17. Lant, C.L.; Roberts, R. Greenbelts in the Cornbelt: Riparian wetlands, intrinsic values, and market failure. *Environ. Plan.* **1990**, *22*, 1375–1388.
18. Termorshuizen, J.W.; Opdam, P.; van den Brink, A. Incorporating ecological sustainability into landscape planning. *Landsc. Urban Plan.* **2007**, *79*, 374–384.
19. Benedict, M.A.; McMahon, E.T. *Green Infrastructure: Linking Landscapes and Communities*; Island Press: Washington, DC, USA, 2006; pp. 57–84.
20. Gómez-Baggethun, E.; Barton, D.N. Classifying and valuing ecosystem services for urban planning. *Ecol. Econ.* **2013**, *86*, 235–245.
21. Haddad, N.M.; Tewksbury, J.J. Low-quality habitat corridors as movement conduits for two butterfly species. *Ecol. Appl.* **2005**, *15*, 250–257.
22. Jongman, R.H.G.; Pungetti, G. *Ecological Networks and Greenways*; Cambridge University Press: Cambridge, MA, USA, 2005.
23. Rapport, D.J.; Costanza, R.; McMichael, A.J. Assessing ecosystem health. *Trends. Ecol. Evol.* **1998**, *13*, 397–402.
24. Bolund, P.; Hunhammar, S. Ecosystem services in urban areas. *Ecol. Econ.* **1999**, *29*, 293–301.
25. Grimm, N.B.; Faeth, S.H.; Golubiewski, N.E.; Redman, C.L.; Wu, J.G.; Bai, X.M.; Briggs, J.M. Global change and the ecology of cities. *Science* **2008**, *319*, 756–760.

26. Niemela, J.; Saarela, S.R.; Soderman, T.; Kopperoinen, L.; Yli-Pelkonen, V.; Vare, S.; Kotze, D.J. Using the ecosystem services approach for better planning and conservation of urban green spaces: A Finland case study. *Biodivers. Conserv.* **2010**, *19*, 3225–3243.
27. Nowak, D.J.; Crane, D.E. Carbon storage and sequestration by urban trees in the USA. *Environ. Pollut.* **2002**, *116*, 381–389.
28. Ayres, R.U.; van den Bergh, J.C.J.M. A theory of economic growth with material/energy resources and dematerialization: Interaction of three growth mechanisms. *Ecol. Econ.* **2005**, *55*, 96–118.
29. Krausmann, F.; Gingrich, S.; Eisenmenger, N.; Erb, K.H.; Haberl, H.; Fischer-Kowalski, M. Growth in global materials use, GDP and population during the 20th century. *Ecol. Econ.* **2009**, *68*, 2696–2705.
30. Fuchs, V.J.; Mihelcic, J.R.; Gierke, J.S. Life cycle assessment of vertical and horizontal flow constructed wetlands for wastewater treatment considering nitrogen and carbon greenhouse gas emissions. *Water Res.* **2011**, *45*, 2073–2081.
31. Ko, J.Y.; Day, J.W.; Lane, R.R.; Day, J.N. A comparative evaluation of money-based and energy-based cost benefit analyses of tertiary municipal wastewater treatment using forested wetlands vs. sand filtration in Louisiana. *Ecol. Econ.* **2004**, *49*, 331–347.
32. Shao, L.; Chen, G.; Hayat, T.; Alsaedi, A. Systems ecological accounting for wastewater treatment engineering: Method, indicator and application, Ecological Indicators. *Ecol. Econ.* **2014**, *47*, 32–42.
33. Zhu, X.D.; Ye, Y.P. Estimate of life-cycle greenhouse gas emissions from a vertical subsurface flow constructed wetland and conventional wastewater treatment plants: A case study in China. *Ecol. Eng.* **2011**, *37*, 248–254.
34. Chen, B.; Chen, Z.M.; Zhou, Y.; Zhou, J.B.; Chen, G.Q. Emergy as embodied energy based assessment for local sustainability of a constructed wetland in Beijing. *Commun. Nonlinear Sci. Numer. Simul.* **2009**, *14*, 622–635.
35. Chen, G.Q.; Shao, L.; Chen, Z.M.; Li, Z.; Zhang, B.; Chen, H.; Wu, Z. Low-carbon assessment for ecological wastewater treatment by a constructed wetland in Beijing, *Ecol. Eng.* **2011**, *37*, 622–628.
36. Chen, Z.M.; Chen, G.Q.; Chen, B.; Zhou, J.B.; Yang, Z.F.; Zhou, Y. Net ecosystem services value of wetland: Environmental economic account. *Commun. Nonlinear Sci. Numer. Simul.* **2009**, *14*, 2837–2843.
37. Forman, R.T.T.; Godron, M. *Landscape Ecology*; John Wiley and Sons, Inc.: New York, NY, USA, 1986; pp. 3–23.
38. Larkin, J.L.; Maehr, D.S.; Hctor, T.S.; Orlando, M.A.; Whitney, K. Landscape linkages and conservation planning for the black bear in west-central Florida. *Anim. Conserv.* **2004**, *7*, 23–34.
39. LaRue, M.A.; Nielsen, C.K. Modelling potential dispersal corridors for cougars in midwestern North America using least-cost path methods. *Ecol. Modell.* **2008**, *212*, 372–381.
40. Meegan, R.P.; Maehr, D.S. Landscape conservation and regional planning for the Florida panther. *Southeast Natl.* **2002**, *1*, 217–232.
41. Penrod, K.; Wildlands, S.C. *South Coast Missing Linkages Project: A Linkage Design for the Santa Monica-Sierra Madre Connection*; South Coast Wildlands: Idyllwild, CA, USA, 2006; pp. 20–52.
42. Schadt, S.; Knauer, F.; Kaczensky, P.; Revilla, E.; Wiegand, T.; Trepl, L. Rule-based assessment of suitable habitat and patch connectivity for the Eurasian lynx. *Ecol. Appl.* **2002**, *12*, 1469–1483.

43. Cook, E.N.; van Lier, H.N. *Landscape Planning and Ecological Networks*; Elsevier Science Ltd: New York, NY, USA, 1994; p. 354.
44. Singleton, P.H.; Gaines, W.L.; Lehmkuhl, J.F. *Landscape Permeability for Large Carnivores in Washington: A Geographic Information System Weighted-Distance and Least-Cost Corridor Assessment*; USDA Forest Service, Pacific Northwest Research Station: Portland, OR, USA, 2002.
45. Anton, C.; Young, J.; Harrison, P.A.; Musche, M.; Bela, G.; Feld, C.K.; Harrington, R.; Haslett, J.R.; Pataki, G.; Rounsevell, M.D. Research needs for incorporating the ecosystem service approach into EU biodiversity conservation policy. *Biodivers. Conserv.* **2010**, *19*, 2979–2994.
46. Costanza, R.; Wilson, M.A.; Troy, A.; Voinov, A.; Liu, S.; D’Agostino, J. The value of New Jersey’s ecosystem services and natural capital. *Gund Institute for Ecological Economics*; University of Vermont and New Jersey Department of Environmental Protection: Trenton, NJ, USA, 2006; pp. 8–52.
47. Hoffmeister, T.S.; Vet, L.E.; Biere, A.; Holsinger, K.; Filser, J. Ecological and evolutionary consequences of biological invasion and habitat fragmentation. *Ecosystems* **2005**, *8*, 657–667.
48. Teng, M.; Wu, C.; Zhou, Z.; Lord, E.; Zheng, Z. Multipurpose greenway planning for changing cities: A framework integrating priorities and a least-cost path model. *Landsc. Urban Plan.* **2011**, *103*, 1–14.
49. Vos, C.C.; Verboom, J.; Opdam, P.F.; Ter Braak, C.J. Toward ecologically scaled landscape indices. *Am. Nat.* **2001**, *157*, 24–41.
50. Corry, R.C.; Laforteza, R.; Brown, R.D. Ecological functionality of landscapes with alternative rehabilitations of depleted aggregate sites. *Int. J. Min. Reclam. Environ.* **2010**, *24*, 216–232.
51. Jesus, R.G.; Gabriel, D.B.; Beatriz, D. Assessing functional landscape connectivity for disturbance propagation on regional scales-A cost-surface model approach applied to surface fire spread. *Ecol. Modell.* **2008**, *211*, 121–141.
52. Weber, T.; Sloan, A.; Wolf, J. Maryland’s green infrastructure assessment: Development of a comprehensive approach to land conservation. *Landsc. Urban Plan.* **2006**, *77*, 94–110.
53. Forester, J.D.; Ives, A.R.; Turner, M.G.; Anderson, D.P.; Fortin, D.; Beyer, H.L.; Smith, D.W.; Boyce, M.S. State-space models link elk movement patterns to landscape characteristics in Yellowstone National Park. *Ecol. Monogr.* **2007**, *77*, 285–299.
54. Schick, R.S.; Loarie, S.R.; Colchero, F.; Best, B.D.; Boustany, A.; Conde, D.A.; Halpin, P.N.; Joppa, L.N.; McClellan, C.M.; Clark, J.S. Understanding movement data and movement processes: Current and emerging directions. *Ecol. Lett.* **2008**, *11*, 1338–1350.
55. Wu, H.; Li, B.; Springer, T.A.; Neill, W.H. Modelling animal movement as a persistent random walk in two dimensions: Expected magnitude of net displacement. *Ecol. Modell.* **2000**, *132*, 115–124.
56. Zetterberg, A.; Mörtberg, U.M.; Balfors, B. Making graph theory operational for landscape ecological assessments, planning, and design. *Landsc. Urban Plan.* **2010**, *95*, 181–191.
57. Zhang, L.; Wang, H. Planning an ecological network of Xiamen Island (China) using landscape metrics and network analysis. *Landsc. Urban Plan.* **2006**, *78*, 449–456.
58. Kong, F.; Yin, H.; Nakagoshi, N.; Zong, Y. Urban green space network development for biodiversity conservation: Identification based on graph theory and gravity modeling. *Landsc. Urban Plan.* **2010**, *95*, 16–27.

59. Turner, M.G. Landscape ecology: The effect of pattern on process. *Annu. Rev. Ecol. Syst.* **1989**, *20*, 171–197.
60. *Arcmap 10.1*; Institute, E.S.R.: Redlands, CA, USA, 2012.
61. De Groot, R.S.; Wilson, M.A.; Boumans, R.M.J. A typology for the classification, description and valuation of ecosystem functions, goods and services. *Ecol. Econ.* **2002**, *41*, 393–408.
62. Troy, A.; Wilson, M.A. Mapping ecosystem services: Practical challenges and opportunities in linking GIS and value transfer. *Ecol. Econ.* **2006**, *60*, 435–449.
63. Millennium Ecosystem Assessment. *Ecosystems and Human Well-Being: Synthesis*; Island Press: Washington, DC, USA, 2005; pp. 39–88.
64. Heal, G.M.; Barbier, E.B.; Boyle, K.J.; Covich, A.P.; Gloss, S.P.; Hershner, C.H.; Hoehn, J.P.; Pringle, C.M.; Polasky, S.; Segerson, K.; *et al.* *Valuing Ecosystem Services: Toward Better Environmental Decision Making*; The National Academies Press: Washington, DC, USA, 2005; pp. 33–58.
65. Villa, F.; Wilson, M.A.; de Groot, R.; Farber, S.; Costanza, R.; Boumans, R.M.J. Designing an integrated knowledge base to support ecosystem services valuation. *Ecol. Econ.* **2002**, *41*, 445–456.
66. Kreuter, U.P.; Harris, H.G.; Matlock, M.D.; Lacey, R.E. Change in ecosystem service values in the San Antonio area, Texas. *Ecol. Econ.* **2001**, *39*, 333–346.
67. Li, T.; Wenkai, L.; Zhenghan, Q. Variations in ecosystem service value in response to land use changes in Shenzhen. *Ecol. Econ.* **2010**, *69*, 1427–1435.
68. Portela, R.; Rademacher, I. A dynamic model of patterns of deforestation and their effect on the ability of the Brazilian Amazonia to provide ecosystem services. *Ecol. Modell.* **2001**, *143*, 115–146.
69. Tong, C.; Feagin, R.A.; Lu, J.; Zhang, X.; Zhu, X.; Wang, W.; He, W. Ecosystem service values and restoration in the urban Sanyang wetland of Wenzhou, China. *Ecol. Eng.* **2007**, *29*, 249–258.
70. Rosenberger, R.S.; Stanley, T.D. Measurement, generalization, and publication: Sources of error in benefit transfers and their management. *Ecol. Econ.* **2006**, *60*, 372–378.
71. Fitter, A.; Elmqvist, T.; Haines-young, R.; Poschin, M.; Rinaldo, A.; Setälä, H.; Stoll-Kleemann, S.; Zobel, M.; Murlis, J. In *Ecosystem Services*; Harrison, R.M., Hester, R.E., Eds.; Royal Society of Chemistry: Cambridge, UK, 2010; pp. 1–28.
72. Liu, S.; Costanza, R.; Troy, A.; D’Aagostino, J.; Mates, W. Valuing New Jersey’s ecosystem services and natural capital: A spatially explicit benefit transfer approach. *Environ. Manag.* **2010**, *45*, 1271–1285.
73. Yang, W.; Chang, J.; Xu, B.; Peng, C.; Ge, Y. Ecosystem service value assessment for constructed wetlands: A case study in Hangzhou, China. *Ecol. Econ.* **2008**, *68*, 116–125.
74. Chen, N.; Li, H.; Wang, L. A GIS-based approach for mapping direct use value of ecosystem services at a county scale: Management implications. *Ecol. Econ.* **2009**, *68*, 2768–2776.
75. Sutton, P.C.; Costanza, R. Global estimates of market and non-market values derived from nighttime satellite imagery, land cover, and ecosystem service valuation. *Ecol. Econ.* **2002**, *41*, 509–527.
76. Turner, R.; Daily, G. The ecosystem services framework and natural capital conservation. *Environ. Resour. Econ.* **2008**, *39*, 25–35.

77. Shrestha, R.K.; Loomis, J.B. Meta-analytic benefit transfer of outdoor recreation economic values: Testing out-of-sample convergent validity. *Environ. Resour. Econ.* **2003**, *25*, 79–100.
78. Bergstrom, J.C.; Taylor, L.O. Using meta-analysis for benefits transfer: Theory and practice. *Ecol. Econ.* **2006**, *60*, 351–360.
79. De Groot, R.; Brander, L.; van der Ploeg, S.; Costanza, R.; Bernard, F.; Braat, L.; Christie, M.; Crossman, N.; Ghermandi, A.; Hein, L.; *et al.* Global estimates of the value of ecosystems and their services in monetary units. *Ecosyst. Serv.* **2012**, *1*, 50–61.
80. MeasuringWorth. Available online: <http://www.measuringworth.com> (accessed on 5 May 2013).
81. Penrod, K.; Cabanero, C.; Beier, P.; Luke, C.; Spencer, W.; Rubin, E.; Paulman, C. *A Linkage Design for the Joshua Tree-Twenty-nine Palms Connection*; South Coast Wildlands: Fair Oaks, CA, USA; Unpublished Report. Available online: <http://www.scwildlands.org> (accessed on 5 March 2013).
82. Finke, J.; Sonnenschein, M. *Information Technologies in Environmental Engineering*; Gomez, J.M., Sonnenschein, M., Muller, M., Welsch, H., Rautenstrauch, C., Eds.; Springer: Berlin, Germany, 2007; pp. 433–444.
83. Tyrväinen, L. Economic valuation of urban forest benefits in Finland. *J. Environ. Manag.* **2001**, *62*, 75–92.
84. Creel, M.; Loomis, J. Recreation value of water to wetlands in the San Joaquin Valley: Linked multinomial logit and count data trip frequency models. *Water Resour. Res.* **1992**, *28*, 2597–2606.
85. Mahan, B.L.; Polasky, S.; Adams, R.M. Valuing urban wetlands: A property price approach. *Land Econ.* **2000**, *76*, 100–113.
86. Mendoza-González, G.; Martínez, M.; Lithgow, D.; Pérez-Maqueo, O.; Simonin, P. Land use change and its effects on the value of ecosystem services along the coast of the Gulf of Mexico. *Ecol. Econ.* **2012**, *82*, 23–32.
87. Natuhara, Y. Ecosystem services by paddy fields as substitutes of natural wetlands in Japan. *Ecol. Eng.* **2013**, *56*, 97–106.
88. Alvarez-Farizo, B. Estimating the benefits of agri-environmental policy: Econometric issues in open-ended contingent valuation studies. *J. Environ. Plan. Manag.* **1999**, *42*, 23–43.
89. Bergstrom, J.C.; Stoll, J.R.; Titre, J.P.; Wright, V.L. Economic value of wetlands-based recreation. *Ecol. Econ.* **1990**, *2*, 129–147.
90. Su, S.; Xiao, R.; Jiang, Z.; Zhang, Y. Characterizing landscape pattern and ecosystem service value changes for urbanization impacts at an eco-regional scale. *Appl. Geogr.* **2012**, *34*, 295–305.
91. Amigues, J.-P.; Desaignes, B.; Gauthier, C.; Keith, J.E. The benefits and costs of riparian analysis habitat preservation: A willingness to accept/willingness to pay contingent valuation approach. *Ecol. Econ.* **2002**, *43*, 17–31.
92. Azar, C.; Sterner, T. Discounting and distributional considerations in the context of global warming. *Ecol. Econ.* **1996**, *19*, 169–184.
93. Bennett, R. The value of footpath provision in the countryside: A case-study of public access to urban-fringe woodland. *J. Environ. Plan. Manag.* **1995**, *38*, 409–418.
94. Bishop, K. Assessing the benefits of community forests: An evaluation of the recreational use benefits of two urban fringe woodlands. *J. Environ. Plan. Manag.* **1992**, *35*, 63–76.
95. Fankhauser, S. The social costs of greenhouse gas emissions: An expected value approach. *Energy J.* **1994**, 157–184.

96. Garrod, G.D.; Willis, K.G. The non-use benefits of enhancing forest biodiversity: A contingent ranking study. *Ecol. Econ.* **1997**, *21*, 45–61.
97. Hope, C.; Maul, P. Valuing the impact of CO₂ emissions. *Energ. Policy* **1996**, *24*, 211–219.
98. Hougner, C.; Colding, J.; Söderqvist, T. Economic valuation of a seed dispersal service in the Stockholm National Urban Park, Sweden. *Ecol. Econ.* **2006**, *59*, 364–374.
99. Maddison, D. A cost-benefit analysis of slowing climate change. *Energ. Policy* **1995**, *23*, 337–346.
100. Newell, R.G.; Pizer, W.A. Discounting the distant future: How much do uncertain rates increase valuations? *J. Environ. Plan. Manag.* **2003**, *46*, 52–71.
101. Nordhaus, W.D.; Popp, D. What is the value of scientific knowledge? An application to global warming using the PRICE model. *Energy J.* **1997**, *18*, 1–46.
102. Plambeck, E.L.; Hope, C. PAGE95: An updated valuation of the impacts of global warming. *Energ. Policy* **1996**, *24*, 783–793.
103. Prince, R.; Ahmed, E. Estimating individual recreation benefits under congestion and uncertainty. *J. Leis. Res.* **1989**, *21*, 61–76.
104. Reilly, J.M.; Richards, K.R. Climate change damage and the trace gas index issue. *Environ. Resour. Econ.* **1993**, *3*, 41–61.
105. Roughgarden, T.; Schneider, S.H. Climate change policy: Quantifying uncertainties for damages and optimal carbon taxes. *Energ. Policy* **1999**, *27*, 415–429.
106. Schauer, M.J. Estimation of the greenhouse gas externality with uncertainty. *Environ. Resour. Econ.* **1995**, *5*, 71–82.
107. Tol, R.S. The marginal costs of greenhouse gas emissions. *Energy J.* **1999**, *20*, 61–81.
108. Willis, K.G.; Garrod, G. An individual travel-cost method of evaluating forest recreation. *J. Agric. Econ.* **1991**, *42*, 33–42.
109. Cordell, H.K.; Bergstrom, J.C. Comparison of recreation use values among alternative reservoir water level management scenarios. *Water Resour. Res.* **1993**, *29*, 247–258.
110. Duffield, J.W.; Neher, C.J.; Brown, T.C. Recreation benefits of instream flow: Application to Montana's Big Hole and Bitterroot Rivers. *Water Resour. Res.* **1992**, *28*, 2169–2181.
111. Kahn, J.R.; Buerger, R.B. Valuation and the consequences of multiple sources of environmental deterioration: The case of the New York striped bass fishery. *J. Environ. Manag.* **1994**, *40*, 257–273.
112. Oster, S. Survey results on the benefits of water pollution abatement in the Merrimack River Basin. *Water Resour. Res.* **1977**, *13*, 882–884.
113. Patrick, R.; Fletcher, J.; Lovejoy, S.; Beek, W.V.; Holloway, G.; Binkley, J. Estimating regional benefits of reducing targeted pollutants: An application to agricultural effects on water quality and the value of recreational fishing. *J. Environ. Manage.* **1991**, *33*, 301–310.
114. Ribaud, M.O.; Epp, D.J. The importance of sample discrimination in using the travel cost method to estimate the benefits of improved water quality. *Land Econ.* **1984**, *60*, 397–403.
115. Sanders, L.D.; Walsh, R.G.; Loomis, J.B. Toward empirical estimation of the total value of protecting rivers. *Water Resour. Res.* **1990**, *26*, 1345–1357.
116. Johnson, K.A.; Polasky, S.; Nelson, E.; Pennington, D. Uncertainty in ecosystem services valuation and implications for assessing land use tradeoffs: An agricultural case study in the Minnesota River Basin. *Ecol. Econ.* **2012**, *79*, 71–79.

117. Ives, C.D.; Hose, G.C.; Nipperess, D.A.; Taylor, M.P. Environmental and landscape factors influencing ant and plant diversity in suburban riparian corridors. *Landsc. Urban Plan.* **2011**, *103*, 372–382.
118. Baschak, L.A.; Brown, R.D. An ecological framework for the planning, design and management of urban river greenways. *Landsc. Urban Plan.* **1995**, *33*, 211–225.
119. Little, C.E. *Greenways for America*; Johns Hopkins University Press: Baltimore, MD, USA, 1995; pp. 81–104.
120. OECD. *Multifunctionality Towards an Analytical Framework*; OECD: Paris, France, 2001; p. 158.
121. Justin, K.; Lant, C.; Shaikh, S.; Wang, G. The geography of ecosystem service value: The case of the Des Plaines and Cache River wetlands, Illinois. *Appl. Geogr.* **2011**, *31*, 303–311.
122. Taylor, P.D.; Fahrig, L.; Henein, K.; Merriam, G. Connectivity is a vital element of landscape structure. *Oikos* **1993**, *68*, 571–573.
123. With, K.A.; Gardner, R.H.; Turner, M.G. Landscape connectivity and population distributions in heterogeneous environments. *Oikos* **1997**, *78*, 151–169.
124. Lee, J.; Kim, J.; Yoo, M.; Kim, E.; Chon, J. Satisfaction and anticipated benefits on a community-based riparian greenway. *Seoul Stud.* **2010**, *11*, 15–28.
125. Wolf, K.L.; Noe, G.B.; Ahn, C. Hydrologic Connectivity to Streams Increases Nitrogen and Phosphorus Inputs and Cycling in Soils of Created and Natural Floodplain Wetlands. *J. Environ. Qual.* **2013**, *42*, 1245–1255.

© 2014 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/4.0/>).